

Symmetries and reductions of order for certain nonlinear third and second-order differential equations with arbitrary nonlinearity

KM Tamizhmani¹, K Krishnakumar¹ and PGL Leach²

¹ Department of Mathematics, Pondicherry University, Kalapet,
Puducherry 605 014, India

²Department of Mathematics, Durban University of Technology, POBox 1334,
Durban 4000, Republic of South Africa.

Abstract

We examine the reductions of the order of certain third- and second-order nonlinear equations with arbitrary nonlinearity through their symmetries and some appropriate transformations. We use the folding transformation which enables one to change from a nonlinearity with an arbitrary exponent to a nonlinearity with a specific numerical exponent.

MSC Numbers: 17B80; 34M15; 58J70

PACS Numbers: 02.20.Sv; 02.30.Hq; 02.30.Ik

Keywords: Folding transformation; Symmetry analysis; Abel's equation;

1 Introduction

Nonlinearity plays an important role in various fields of mathematics, physics, biology *etc.* The determination of closed-form solution of nonlinear ordinary differential equations is quite difficult in general. Around the turn of twentieth century Painlevé and his school classified second-order nonlinear ordinary differential equations based on what is known as the Painlevé Property. An ordinary differential equation is said to possess the Painlevé Property if all solutions are single-valued around all movable singularities. They arrived at fifty canonical equations of which the solutions of six equations could not be expressed in terms of then known functions. These six equations are the famous Painlevé equations [1]. There exists another class of equations which are linearizable through transformations of dependent/or independent variable. The famous equation among this class is the Gambier system which is written in the form of coupled Riccati equations in cascade. The Gambier equation can be rewritten as second-order nonlinear ordinary differential equation of the form [2]

$$y'' = \left(I - \frac{I}{n}\right) \frac{y'^2}{y} + \left(ay + b + \frac{c}{y}\right) y' - \frac{na^2}{(n+2)^2} y^3 + ey^2 + fy + g + \frac{h}{y}, \quad (1.1)$$

where here and below, unless otherwise indicated, the prime denotes differentiation with respect to the independent variable, x . For each n one can obtain a condition on a, b and c which guarantees the Painlevé Property (*ie* absence of movable critical points). There were detailed analyses to determine integrable/linearizable nonlinear ordinary differential equations by Painlevé, Gambier, Garnier, Bureau [3–9] and others through the analysis of their singularity structures. In another development Sophus Lie formulated criteria for linearising certain classes of nonlinear second-order differential equations through the approach of symmetry [10]. Later Chazy [11] gave conditions for linearisation and integration of certain higher-order nonlinear ordinary differential equations. Many authors have been using various methods such as symmetry analysis, the method of the Jacobi Last Multiplier [12–18], the Prelle-Singer method [23–26] *etc* to find the solutions/first integrals of nonlinear ordinary differential equations. Among these methods that of symmetry analysis has been used by researchers effectively [19–21]. Recently finding the interconnection among these methods has also been a very interesting part of research [27, 28].

Our main objective of this paper is to investigate the third-order nonlinear ordinary differential equation of Chazy [11] through symmetry analysis. From the point of view of symmetry we look for reductions of this equation to well-known equations. We show that one of the reductions of the Chazy equation is an Abel equation of the second kind. Our results motivated us to reexamine the list of new second-order nonlinear ordinary differential equations given in a series of papers [24, 26]. We show through the approach of symmetry and the implementation of a series of transformations that all these equations can be transformed to well-known equations which are found in the literature. We compute the Lie point symmetries of these equations by using the Mathematica add-on, Sym [29–32]. One can reduce the order of the given differential equations by using canonical variables which are found from the symmetries of the differential equations. Based on this idea we firstly give the reduction of a third-order ordinary differential equation which was used by Chazy [11]. Although he reduced the third-order equation to an Abel equation of the first kind, we reduce it to an Abel equation of the second kind¹. Although the Abel equation of the second kind is not integrable for arbitrary parameters, there exist particular solutions of them with certain restrictions on the parameters. These restrictions are already given in [33].

Secondly we examine the general form of certain second-order nonlinear

¹The two are kinds of Abel's equations are related by a transformation.

equations,

$$y'' + (k_1 y^q + k_2) y' + k_3 y^{2q+1} + k_4 y^{q+1} + k_5 y = 0 \quad (1.2)$$

and

$$y'' + \lambda \frac{y'^2}{y} + (k_1 y + k_2) y' + K_3 y^3 + K_4 y^2 + K_5 y = 0, \quad (1.3)$$

where $q, \lambda, k_1, k_2, k_3, k_4, k_5, K_3, K_4$ and K_5 are arbitrary parameters. For particular values of the parameters in these equations one can obtain many well-known equations of oscillators such as the modified Emden equation, Emden equation *etc.* [23]. Eq.(1.2) and eq.(1.3) have already been analyzed [24,26] separately through the extended Prelle-Singer method. By this method the authors of [24,26] claimed that the following equations with some restrictions on the parameters are new integrable equations. The first five equations (1.4–1.8) are from [24]. The remaining (1.9–1.12) are from [26]. The equations are

$$y'' + k_1 y y' + k_3 y^3 + k_5 y = 0, \quad (1.4)$$

$$y'' + (k_1 y + k_2) y' + k_3 y^3 + \frac{k_1 k_2}{3} y^2 + \frac{2k_2^2}{9} y = 0, \quad (1.5)$$

where $k_3 = \frac{(r-1)k_1^2}{2r^2}$.

$$y'' + (k_1 y^2 + k_2) y' + k_3 y^5 + \frac{4(r-1)k_1 k_2}{3r^2} y^3 + \frac{(r-1)k_2^2}{r^2} y = 0, \quad (1.6)$$

$$y'' + (k_1 y^2 + k_2) y' + \frac{k_1^2}{16} y^5 + \frac{k_1 k_2}{4} y^3 + k_5 y = 0, \quad (1.7)$$

$$y'' + (k_1 y^2 + k_2) y' + k_3 y^5 + \frac{k_1 k_2}{4} y^3 + \frac{3k_2^2}{16} y = 0, \quad (1.8)$$

where $k_3 = \frac{(r-1)k_1^2}{3r^2}$ and r is an arbitrary parameter.

$$y'' + \lambda \frac{y'^2}{y} + (k_1 y + k_2) y' + K_3 y^3 + \frac{k_2 K_3 (3 + 2\lambda)}{k_1 (1 + \lambda)} y^2 + \frac{K_3 k_2^2 (2 + \lambda)}{k_1^2 (1 + \lambda)} y = 0, \quad (1.9)$$

$$y'' + \lambda \frac{y'^2}{y} + (k_1 y + k_2) y' + \frac{(1 + \lambda) k_1^2}{(3 + 2\lambda)^2} y^3 + \frac{k_1 k_2}{(3 + 2\lambda)} y^2 + K_5 y = 0, \quad (1.10)$$

$$y'' + \lambda \frac{y'^2}{y} + k_2 y' + K_4 y^2 + \frac{2(3 + 2\lambda) k_2^2}{(5 + 4\lambda)^2} y = 0, \quad (1.11)$$

$$y'' + \lambda \frac{y'^2}{y} + (k_1 y + k_2) y' + K_3 y^3 + \frac{k_1 k_2}{(3 + 2\lambda)} y^2 + \frac{(2 + \lambda) k_2^2}{(3 + 2\lambda)^2} y = 0. \quad (1.12)$$

Our investigation reveals that all the above new integrable equations can be identified with well-known equations, such as the Riccati equation, Bernoulli equation, Abel equations of the first and second kinds and the Emden-Fowler equation, by taking a series of transformations that are already given in [33]. As the determination of these solutions and conditions for linearisation of equations of these types are well-known, we can show that the equations listed above are not essentially new equations but variations on the types mentioned above. For both third- and second-order equations we use appropriate folding transformations² which takes arbitrary exponents to a specific numerical nonlinearity. Otherwise finding the symmetries of these equations with arbitrary nonlinearity is difficult.

2 Symmetries and reductions of the third-order equation

We consider equation of the form [2]

$$-y''' + Ay^q y'' + By^{q-1} y'^2 + Fy^{2q} y' + Gy^{3q+1} = 0. \quad (2.1)$$

When one applies the folding transformation, $v(x) = y^{-q}$, one gets the form

$$\begin{aligned} v^2 v^{(3)} - 3 \left(1 + \frac{1}{q}\right) v v' v'' - A v v'' + \left(2 + \frac{1}{q^2} + \frac{3}{q}\right) v'^3 \\ + \left(A + \frac{A}{q} + \frac{B}{q}\right) v'^2 - F v' + G q = 0. \end{aligned} \quad (2.2)$$

Now the arbitrary nonlinearity of (2.1) has been removed by the above folding transformation. The symmetries of (2.2) are

$$\Gamma_1 = \partial_x \quad (2.3)$$

$$\Gamma_2 = x \partial_x + v \partial_v. \quad (2.4)$$

According to the symmetry Γ_1 the canonical variables are $(r = v, R = v')$. By using these canonical variables one can find the transformation

$$v = r, \quad v' = R(r)$$

²Okamoto *et al* [34] introduced the folding transformation. This is a class of transformations of algebraic rather than rational form. This type of transformation plays a crucial role in many of continuous and discrete integrable systems [35, 36]

to reduce the third-order nonlinear ordinary differential equation, (2.2), to a second-order nonautonomous nonlinear ordinary differential equation given by

$$q^2 r^2 (RR'^2 + R^2 R'') - 3qr(1+q)R^2 R' - Aq^2 r R R' + (1+3q+2q^2)R^3 + (Aq+Bq+Aq^2)R^2 - Fq^2 R + Gq^3 = 0, \quad (2.5)$$

where here ' represents differentiation with respect to r . The symmetry of (2.5) is given by

$$\Gamma_1 = r\partial_r, \quad (2.6)$$

and corresponding canonical variables are ($z = \log r$, $Z = R$). Thereby we obtain the transformation

$$R = Z, \quad rR' = Z'$$

and the second-order autonomous ordinary differential equation

$$qZ^2 (q(Z'' - Z') - 3(1+q)Z' + Aq + A + B) + (1+3q+2q^2)Z^3 - q^2 Z (F + AZ' - Z'^2) + Gq^3 = 0, \quad (2.7)$$

where now ' represents differentiation with respect to z . The symmetry of (2.7) is given by

$$\Gamma_1 = \partial_z. \quad (2.8)$$

The symmetry leads to the canonical variables ($h = Z$, $z = T$) and the transformations

$$Z = h, \quad Z' = H$$

which reduce equation (2.7) to an Abel's equation of the second kind,

$$HH' + \frac{H^2}{h} - \frac{AH}{h} - \left(4 + \frac{3}{q}\right)H + \frac{Gq}{h^2} - \frac{F}{h} + 2h + \frac{h}{q^2} + \frac{A}{q} + \frac{B}{q} + \frac{3h}{q} + A = 0, \quad (2.9)$$

where ' represents differentiation with respect to h . The Abel's equation of the second kind (2.9) is not integrable for arbitrary parameters. However, we give particular solutions of this equation with certain possible restrictions among the parameters. The details can be found in [33]. For the sake of completeness we reproduce the results given in [33] for the various cases of (2.9) listed below. Here we excluded the trivial linear cases.

Case 1:

$$A = \text{arbitrary}, B = -\frac{A}{4}, F = \text{arbitrary}, G = 0, q = -\frac{3}{4}$$

$$HH' + \frac{H^2}{h} - \frac{AH}{h} - \frac{2h}{9} - \frac{F}{h} = 0. \quad (2.10)$$

The transformation $H = A \frac{w(h)}{h}$ [see [33], 1.3.4-2, equation 6] reduces this to

$$ww' - w - \frac{2h^3}{9A^2} - \frac{Fh}{A^2} = 0. \quad (2.11)$$

One has a form of the Emden-Fowler equation

$$\phi'' = C\xi^n\phi^3 \quad (2.12)$$

from (2.11) by the transformation [see [33], 1.3.1-2 equation 71]

$$h = \phi\xi^{1+\frac{n}{2}}, w = \xi^{1+\frac{n}{2}} \frac{(2+n)\phi + 2\xi\phi'}{2(3+n)},$$

where

$$\frac{F}{A^2} = -\frac{(2+n)(4+n)}{4(3+n)^2} \Leftrightarrow n = \pm \left(\frac{A}{\sqrt{A^2 + 4F}} - 3 \right),$$

the prime denotes differentiation with respect to ξ and $C = -2(3+n)^2/(9A^2)$.

Subcase: $n = -6$

The solution of (2.12) in parametric form is

$$\xi = aI_1^2 \left[\int (1 \pm \tau^4)^{\frac{-1}{2}} d\tau + I_2 \right]^{-1}, \quad \phi = bI_1^4 \tau \left[\int (1 \pm \tau^4)^{\frac{-1}{2}} d\tau + I_2 \right]^{-1}, \quad (2.13)$$

where $C = \pm 2a^4b^{-2}$.

This result is a consequence of the property that for $n = -6$ (2.12) has the two Lie point symmetries,

$$\Gamma_1 = \xi\partial_\xi + 2\phi\partial_\phi \quad \text{and} \quad \Gamma_2 = \xi^2\partial_\xi + \xi\phi\partial_\phi,$$

whereas in general there is just the single symmetry $\Gamma = 2\xi\partial_\xi - (2+n)\phi\partial_\phi$.

Case 2:

$$A = -2B, B = \text{arbitrary}, F = \text{arbitrary}, G = 0, q = -\frac{3}{2}$$

$$HH' + \frac{H^2}{h} - \frac{2BH}{h} - 2H + \frac{4h}{9} - \frac{F}{h} - \frac{4B}{3} = 0. \quad (2.14)$$

The transformation $H = w(h) + \frac{2}{3}h$ [see [33], 1.3.4-2, equation 7] reduces this equation to

$$(F - 2Bw - w^2) \frac{dh}{dw} = wh + \frac{2}{3}h^2 \quad (2.15)$$

with the solution

$$h = -\frac{3 \exp(L \tan^{-1} M) F}{4B \exp(L \tan^{-1} M) + 2 \exp(L \tan^{-1} M) w - 3FNI}, \quad (2.16)$$

where I is a constant of integration,

$$L = \frac{B}{\sqrt{-B^2 - F}}, \quad M = \frac{(B + w)}{\sqrt{-B^2 - F}} \quad \text{and} \quad N = \sqrt{-F + 2Bw + w^2}.$$

Case 3:

$$A = \frac{5}{2}, B = \text{arbitrary}, F = -\frac{3}{2}, G = 0, q = 1$$

$$HH' + \frac{H^2}{h} - \frac{5H}{2h} - 7H + 6h + \frac{3}{2h} + B + 5 = 0. \quad (2.17)$$

The transformation $h = \xi^2$ and $H = \xi w(\xi) + 2\xi^2 + 1$ [see [33], 1.3.4-2, equation 8] reduces this equation to

$$(2 - 2B - 3w^2) \frac{d\xi}{dw} = 1 + w\xi + 2\xi^2 \quad (2.18)$$

with solution

$$\begin{aligned} \xi = & \left(-24\sqrt{-1 + Bw} I \ P[l, m, n] + \sqrt{11 - 10B - B^2} I \ P[7l, m, n] \right. \\ & - 24\sqrt{-1 + Bw} \ Q[l, m, n] - 7\sqrt{6}\sqrt{-(-1 + B)^2} (1 + \ Q[7l, m, n]) \\ & \left. + \sqrt{6}\sqrt{11 - 10B - B^2} \ Q[7l, m, n] \right) / \\ & \left(12\sqrt{-1 + B} I \ P[l, m, n] + 12\sqrt{-1 + B} \ Q[l, m, n] \right), \end{aligned} \quad (2.19)$$

where P and Q are Legendre functions, I is a constant of integration,

$$l = \frac{1}{6}, \quad m = \frac{1}{6} \sqrt{\frac{(11+B)}{(-1+B)}} \quad \text{and} \quad n = -\frac{\sqrt{3}w}{\sqrt{2(1-B)}}.$$

Case 4:

$$A = \frac{5}{2}, \quad B = \text{arbitrary}, \quad F = -\frac{3}{2}, \quad G = 0, \quad q = -\frac{5}{2}$$

$$HH' + \frac{H^2}{h} - \frac{5H}{2h} - \frac{14H}{5} + \frac{24h}{25} + \frac{3}{2h} - \frac{2B}{5} + \frac{3}{2} = 0. \quad (2.20)$$

The transformation $h = \xi^2$ and $H = \xi w(\xi) + \frac{4}{5}\xi^2 + 1$ [see [33], 1.3.4-2, equation 8] reduces this to

$$(9 + 4B - 15w^2) \frac{d\xi}{dw} = 5 + 5w\xi + 4\xi^2 \quad (2.21)$$

with the solution

$$\begin{aligned} \xi = & - (120wI \ P[i, j, k] + 120w \ Q[i, j, k] \\ & + \sqrt{15}(\sqrt{-39+4B} - 7\sqrt{9+4B})I \ P[i, j, k] \\ & - 7\sqrt{15}\sqrt{9+4B} \ Q[7i, j, k] + \sqrt{-585+60B} \ Q[7i, j, k]) / \\ & (24I \ P[i, j, k] + 24 \ Q[i, j, k]), \end{aligned} \quad (2.22)$$

where P and Q are Legendre functions, I is a constant of integration,

$$i = \frac{1}{6}, \quad j = \frac{1}{6} \sqrt{\frac{(-39+4B)}{(9+4B)}} \quad \text{and} \quad k = \frac{\sqrt{15}w}{\sqrt{9+4B}}.$$

Case 5:

$$A = 0, \quad B = 0, \quad F = \text{arbitrary}, \quad G = 0, \quad q = -\frac{3}{4}$$

$$HH' + \frac{H^2}{h} - \frac{2h}{9} - \frac{F}{h} = 0, \quad (2.23)$$

the solution of which is given by

$$H = \pm \frac{\sqrt{9Fh^2 + h^4 + 9I}}{3h}. \quad (2.24)$$

Case 6:

$$A = -2B, B = \text{arbitrary}, F = 0, G = 0, q = -\frac{1}{2}$$

$$H' + \frac{H}{h} + \frac{2B}{h} + 2 = 0 \quad (2.25)$$

with solution given by [see [33], 1.3.4-2, equation 34]

$$H = -2B - h + \frac{I}{h}. \quad (2.26)$$

Case 7:

$$A = \text{arbitrary}, B = 0, F = 0, G = 0, q = -1$$

$$H' + \frac{H}{h} - \frac{A}{h} - 1 = 0. \quad (2.27)$$

This equation has the solution [see [33], 1.3.4-2, equation 34]

$$H = A + \frac{h}{2} + \frac{I}{h}. \quad (2.28)$$

3 Symmetries and reductions of the second-order equation

We consider the equation (1.2),

$$y'' + (k_1 y^q + k_2) y' + k_3 y^{2q+1} + k_4 y^{q+1} + k_5 y = 0.$$

When we apply the folding transformation, $w = y^q$, we obtain the following form

$$w'' + \left(\frac{1}{q} - 1 \right) \frac{w'^2}{w} + (k_1 w + k_2) w' + q k_3 w^3 + q k_4 w^2 + q k_5 w = 0. \quad (3.1)$$

Remark:

When $(\frac{1}{q} - 1) = \lambda$, $\frac{k_3}{1+\lambda} = K_3$, $\frac{k_4}{1+\lambda} = K_4$ and $\frac{k_5}{1+\lambda} = K_5$, the equation above, (3.1), is equation (1.3). Therefore there is no need to make a separate analysis.

Now we apply the symmetry method to find the further transformations which reduce the order of equation (3.1). The determining equations of (3.1) have four arbitrary functions and conditions over them. These functions do exist in the coefficient functions, ξ and η , according to

$$\xi(x, w) = a(x) + qw^{\frac{1}{q}}b(x) \quad (3.2)$$

$$\begin{aligned} \eta(x, w) = \frac{w^{1-\frac{1}{q}}}{(1+q)(2+q)} & \left[\left(-q^2 w^{\frac{2}{q}} b(x) (2wk_1 + (2+3q+q^2)k_2) \right. \right. \\ & \left. \left. + (2+3q+q^2)(w^{\frac{1}{q}}d(x) + c(x) + q^2 w^{\frac{2}{q}}b'(x)) \right) \right], \end{aligned} \quad (3.3)$$

where $a(x), b(x), c(x)$ and $d(x)$ are arbitrary functions. According to the conditions on the arbitrary functions, there do occur a few cases

Case 1:

$$a(x) = A_0, b(x) = 0, c(x) = 0 \text{ and } d(x) = 0$$

The corresponding symmetry is

$$\Gamma_1 = \partial_x$$

Case 2:

$$a(x) = A_1 - \frac{(2+q)A_0}{qk_2} \exp \left[\frac{qk_2x}{2+q} \right], \quad b(x) = 0, c(x) = 0 \quad \text{and} \quad d(x) = A_0 \exp \left[\frac{qk_2x}{2+q} \right]$$

with the conditions $k_4 = k_1k_2/(2+q)$ and $k_5 = (1+q)k_2^2/(2+q)^2$.

The corresponding symmetries are

$$\Gamma_1 = \partial_x \quad (3.4)$$

$$\Gamma_2 = -\exp \left[\frac{qk_2x}{2+q} \right] \left(\frac{(2+q)}{qk_2} \partial_x + w \partial_w \right). \quad (3.5)$$

Case 3:

$$k_1 = 0, k_3 = 0 \text{ and } k_4 = 0.$$

The corresponding symmetries are

$$\begin{aligned}\Gamma_1 &= w\partial_w, \\ \Gamma_2 &= \partial_x - \frac{1}{2}k_2qw\partial_w, \\ \Gamma_3 &= \exp\left[-\frac{(k_2 + N)x}{2}\right] w^{\left(\frac{-1+q}{q}\right)}\partial_w, \\ \Gamma_4 &= \exp\left[\frac{(-k_2 + N)x}{2}\right] w^{\left(\frac{-1+q}{q}\right)}\partial_w, \\ \Gamma_5 &= \exp[Nx]\frac{(2\partial_x - k_2qw\partial_w + qNw\partial_w)}{2N}, \\ \Gamma_6 &= \exp[-Nx]\frac{(-2\partial_x + k_2qw\partial_w + qNw\partial_w)}{2N}, \\ \Gamma_7 &= \frac{qw^{1/q}}{2} \exp\left[\frac{(k_2 + N)x}{2}\right] (2\partial_x - k_2qw\partial_w + qNw\partial_w) \quad \text{and} \\ \Gamma_8 &= -\frac{qw^{1/q}}{2} \exp\left[\frac{(k_2 - N)x}{2}\right] (-2\partial_x + k_2qw\partial_w + qNw\partial_w),\end{aligned}$$

where $N = \sqrt{k_2^2 - 4k_5}$. When we treat Case 1, we arrive at the conditions $k_4 = k_1k_2/(2+q)$ and $k_5 = (1+q)k_2^2/(2+q)^2$ and the last case contains the conditions $k_1 = 0, k_3 = 0$ and $k_4 = 0$ which means that the given second-order equation is a linear equation. Hence only one reduction is sufficient for all cases by using $\Gamma_1 = \partial_x$.

According to the symmetry Γ_1 the canonical variables are $(r = w, s = x)$. By using these canonical variables one can find the transformation

$$w = r, \quad w' = H(r)$$

to reduce the second-order nonlinear ordinary differential equation, (3.1), to a first-order nonautonomous nonlinear ordinary differential equation with the form of an Abel's equation of the second kind. It is

$$HH' + \left(\frac{1}{q} - 1\right) \frac{H^2}{r} + (k_1r + k_2)H + qk_3r^3 + qk_4r^2 + qk_5r = 0, \quad (3.6)$$

where ' represents differentiation with respect to r . The transformation $H(r) = R(r)r^{1-1/q}$ changes the Abel equation of the second kind into one of the same kind with new dependent variable $R(r)$. It has the form

$$RR' + (k_1r + k_2)r^{\frac{1}{q}-1}R + (qk_3r^2 + qk_4r + qk_5)r^{\frac{2}{q}-1} = 0. \quad (3.7)$$

3.1 Reduction to an Abel equation

If one takes $k_2 = 0$, $k_4 = 0$ and $q = 1$, then (3.7) reduces to an equation of the form

$$RR' + k_1rR + (k_3r^2 + k_5)r = 0. \quad (3.8)$$

We now make the transformation $z = -\frac{1}{2}k_1r^2$ [see [33], 0.1.6-2]. Eq.(3.8) reduces to a solvable Abel equation of the form [see [33], 1.3.1-2, equation 2]

$$R\frac{dR}{dz} - R = \frac{k_5}{k_1} - \frac{2k_3}{k_1^2}z \quad (3.9)$$

with no constraints upon the values of k_1 , k_3 and k_5 ³. The solution in parametric form is

$$z = c \exp \left[- \int \frac{\tau}{\tau^2 - \tau + \frac{2k_3}{k_1^2}} d\tau + \frac{k_1k_5}{2k_3} \right], \quad (3.10)$$

$$R = c \tau \exp \left[- \int \frac{\tau}{\tau^2 - \tau + \frac{2k_3}{k_1^2}} d\tau \right]. \quad (3.11)$$

When one uses (3.10) and (3.11), one can get the solution of (1.4) in the list of the new integrable cases.

3.2 Reduction to a linear first-order equation

If we make the transformation

$$R = r^{\frac{1}{q}+1} \left(t - \frac{qk_2}{(2+q)r} \right),$$

(3.7) reduces to a solvable linear first-order equation of the form [see [33], 1.3.3-2, equation 13 with $a = 0$, $n = 1 + 1/q$, $c = -k_1$, $b = -qk_2/(q+2)$, $d = qk_3$]

$$\left(\left(\frac{1}{q} + 1 \right) t^2 + k_1t + qk_3 \right) \frac{dr}{dt} = \frac{qk_2}{(2+q)} - tr \quad (3.12)$$

³As these parameter occur in ratios, any two are independent.

with conditions on the parameters

$$k_3 - \text{arbitrary}, \quad k_4 = \frac{k_1 k_2}{2+q} \quad \text{and} \quad k_5 = \frac{(1+q)k_2^2}{(2+q)^2}.$$

The solution of (3.12) is [see [33], 0.1.2-5]

$$r = \exp \left(- \int \frac{t dt}{f(t)} \right) \left[\int \left(\exp \int \frac{t dt}{f(t)} \right) \left(\frac{q k_2}{(2+q)f(t)} \right) dt + C \right], \quad (3.13)$$

where $f(t) = (1 + 1/q + 1)t^2 + k_1 t + q k_3$. We now show that, given the results presented above, one can precisely obtain the results given in [24, 26] for various values of q .

For example take $q = 1$. Then (3.13) gives the solution of (1.5) with an arbitrary value of k_3 , $k_4 = \frac{k_1 k_2}{3}$ and $k_5 = \frac{2k_2^2}{9}$ [24]. For $q = 2$ (3.13) gives the solution of (1.8) with an arbitrary value of k_3 , $k_4 = \frac{k_1 k_2}{4}$ and $k_5 = \frac{3k_2^2}{19}$ [24]. If $q = \frac{1}{1+\lambda}$, then (3.13) gives the solution of (1.12) with arbitrary value of K_3 , $K_4 = \frac{k_1 k_2}{(3+2\lambda)}$ and $K_5 = \frac{(2+\lambda)k_2^2}{(3+2\lambda)^2}$ [26].

3.3 Reduction to a solvable Abel equation

If one takes the transformation

$$\xi = -k_1 r^{1/q} \left(\frac{qr}{1+q} + \frac{qk_2}{k_1} \right),$$

eq.(3.7) is reduced to a solvable Abel's equation of the form [see [33], 1.3.3-2, equation 9 with $a = -k_1$, $n = 1/q$, $b = -qk_2/k_1$]

$$R \frac{dR}{d\xi} = R - \frac{(1+q)k_3}{k_1^2} \xi \quad (3.14)$$

with conditions k_3 -arbitrary, $k_4 = \frac{(2+q)k_2 k_3}{k_1}$ and $k_5 = \frac{(1+q)k_2^2 k_3}{k_1^2}$. The solution of (3.14) in parametric form is [see [33], 1.3.1-2, equation 2]

$$\xi = C \exp \left(- \int \frac{\tau d\tau}{\tau^2 - \tau + \frac{(1+q)k_3}{k_1^2}} \right), \quad (3.15)$$

$$R = C\tau \exp\left(-\int \frac{\tau d\tau}{\tau^2 - \tau + \frac{(1+q)k_3}{k_1^2}}\right). \quad (3.16)$$

Take $q = 2$ and $k_3 = (r-1)k_1^2/(3r^2)$. Then (3.16) gives the solution of (1.6) with the conditions $k_4 = \frac{4(r-1)k_1k_2}{3r^2}$ and $k_5 = \frac{(r-1)k_2^2}{r^2}$ [24].

Take $q = \frac{1}{1+\lambda}$. Then (3.16) gives the solution of (1.9) with arbitrary value of K_3 , $K_4 = \frac{k_2K_3(3+2\lambda)}{k_1(1+\lambda)}$ and $K_5 = \frac{K_3k_2^2(2+\lambda)}{k_1^2(1+\lambda)}$ [26]. However, we have obtained solutions (3.15) and (3.16) for arbitrary k_3 .

3.4 Reduction to a Bernoulli equation

If one uses the transformation

$$R = r^{\frac{1}{q}} \left(\phi - \frac{qk_1}{2+q} \right),$$

eq.(3.7) is reduced to a Bernoulli equation of the form [see [33], 1.3.3-2, equation 10 with $a = -\frac{qk_1}{(2+q)}$, $n = \frac{1}{q}$, $b = -k_2$],

$$\left(\frac{\phi^2}{q} + k_2\phi + qk_5 \right) \frac{dr}{d\phi} = -\phi r + \frac{qk_1}{2+q}r^2 \quad (3.17)$$

with conditions $k_3 = \frac{k_1^2}{(2+q)^2}$, $k_4 = \frac{k_1k_2}{(2+q)}$ and k_5 -arbitrary. The solution of (3.17) is [see [33], 0.1.2-6]

$$r^{-1} = \exp\left(-\int \frac{\phi d\phi}{g(\phi)}\right) \left[C - \int \exp\left(\int \frac{\phi d\phi}{g(\phi)}\right) \frac{qk_1}{2+q} d\phi \right], \quad (3.18)$$

where $g(\phi) = \frac{\phi^2}{q} + k_2\phi + qk_5$.

We recover the conditions given in [24] by taking $q = 2$ from (3.7). Then

(3.18) gives the solution of (1.7) with an arbitrary value of k_5 and the conditions $k_3 = \frac{k_1^2}{16}$ and $k_4 = \frac{k_1k_2}{4}$ [24]. Take $q = \frac{1}{1+\lambda}$. Then (3.18) gives

the solution of (1.10) with an arbitrary value of K_5 , $K_3 = \frac{(1+\lambda)k_1^2}{(3+2\lambda)^2}$ and $K_4 = \frac{k_1k_2}{(3+2\lambda)}$ [26].

3.5 Reduction to a Riccati equation

If one takes the transformation

$$\chi = \sqrt{r} \text{ and } R = r^{\frac{2+q}{2q}} \left(t - \frac{2qk_1\sqrt{r}}{4+3q} - \frac{2qk_2}{(4+q)\sqrt{r}} \right),$$

eq.(3.7) is reduced to a Riccati equation of the form [see [33], 1.3.3-2, equation 11],

$$\left[\left(\frac{2+q}{2q} \right) t^2 + c_0 \right] \frac{d\chi}{dt} = \frac{qk_1}{4+3q} \chi^2 - \frac{t}{2} \chi + \frac{qk_2}{4+q}, \quad (3.19)$$

where

$$c_0 = qk_4 - \frac{4q(2+q)k_1k_2}{(4+q)(4+3q)}$$

with conditions

$$k_3 = \frac{2(2+q)k_1^2}{(4+3q)^2}, \quad k_4 \text{-arbitrary and } k_5 = \frac{2(2+q)k_2^2}{(4+q)^2}.$$

The Cole-Hopf transformation,

$$\chi = \frac{-\left(\frac{2+q}{2q}t^2 + c_0\right)\mu'}{\frac{qk_1}{4+3q}\mu},$$

converts the nonlinear Riccati equation (3.19) into a linear second-order equation of the form:

$$\left(\frac{2+q}{2q}t^2 + c_0 \right)^2 \mu''_{tt} + \left(\frac{2+q}{q} + \frac{1}{2} \right) t \left(\frac{2+q}{2q}t^2 + c_0 \right) \mu'_t + \frac{q^2k_1k_2}{(4+q)(4+3q)} \mu = 0. \quad (3.20)$$

Take $q = \frac{1}{1+\lambda}$ and $k_1 = 0$. Then the solution of (3.20) gives the solution of

$$(1.11) \text{ with arbitrary value of } K_4 \text{ and } K_5 = \frac{2(3+2\lambda)k_2^2}{(5+4\lambda)^2} \text{ [26].}$$

4 Conclusion

We have investigated the linearisation of two classes of nonlinear equations, one class being of the third order and the other of the second order. These equations contain parameters. By the performance of a symmetry analysis we have been able to identify those values of the parameters which permit the existence of Lie point symmetries and consequently reduction of order. In particular we were able to identify third-order equations with two Lie point symmetries and this led to reduction to first-order equations generally with the form of an Abel's equation of the second kind. Integrable cases of this class of equations are provided in the compendium of Polyanin and Zeitev [33].

In the case of the second-order equations we were able to produce a uniform examination of equations reported in earlier literature [23–26] as being new additions to the pantheon of integrable second-order equations. Whilst it is true that these equations were of new appearance, we showed that they could be transformed into equations of known integrability. However, there is no denying that they do provide new expressions for integrable equations.

Acknowledgements

KMT would like to thank Professor Tudor Ratiu, CIB, EPFL for the invitation and hospitality where the work was initiated. KK thanks the University Grants Commission for providing a UGC-Basic Scientific Research Fellowship to perform this research work. PGLL thanks Professor KM Tamizhmani and the Department of Mathematics, Pondicherry University, for the invitation and provision of facilities while this work was undertaken.

References

- [1] Ince EL. 1927 *Ordinary Differential Equations*. (Longmans, Green & Co, London).
- [2] Gambier B. 1909 Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes. *Acta. Math.* **33**, 1–55.
- [3] Painlevé P. 1973 *Leçons sur la théorie analytique des équations différentielles*. (Leçons de Stockholm, 1895) (Hermann, Paris, 1897). Reprinted, *Oeuvres de Paul Painlevé*, vol. I, Éditions du CNRS, Paris.

- [4] Painlevé P. 1900 Mémoire sur les équations différentielles du second ordre dont l'intégrale générale est uniforme. *Bull. Math. Soc. France* **28**, 201-265.
- [5] Painlevé P. 1902 Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme. *Acta. Math.* **25**, 1-85.
- [6] Painlevé P. 1906 Sur les équations différentielles du second ordre à points critiques fixes. *C. R. Acad. Sc. Paris* **143**, 1111-1117.
- [7] Garnier R. 1912 Sur les équations différentielles du troisième ordre dont l'intégrale générale est uniforme et sur une classe d'équations nouvelles d'ordre supérieur dont l'intégrale générale a ses points critiques fixes. *Ann sci École norm sup* **XXIX**, 1-126.
- [8] Garnier R. 1960 Sur des systèmes différentiels du second ordre dont l'intégrale est uniforme. *Ann sci École norm sup* **77**, 123-144.
- [9] Bureau FJ. 1964 Differential equations with fixed critical points. *Annali di Matematica pura ed applicata* **64**, 229-364.
- [10] Lie S. 1874 Über die Integration durch bestimmte Integrale von einer Klasse linear partieller Differentialgleichungen. *Math. Ann.* **8**, 328-368.
- [11] Chazy J. 1910, 1911 Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'intégrale générale a ses points critiques fixes. *Acta. Math.* **34**, 317-385.
- [12] Nucci MC, Leach PGL. 2002 Jacobi's last multiplier and the complete symmetry group of the Euler-Poinsot system. *J. Nonlinear Math. Phys.* **9(sup 2)**, 110-121. (doi:10.2991/jnmp.2002.9.s2.10)
- [13] Nucci MC, Leach PGL. 2004 Jacobi's last multiplier and symmetries for the Kepler Problem plus a lineal story. *J. Phys. A: Math. Gen.* **37(31)**, 7743-7753. (doi:10.1088/0305-4470/37/31/007)
- [14] Nucci MC. 2005 Jacobi last multiplier and Lie symmetries: a novel application of an old relationship. *J. Nonlinear Math. Phys.* **12(2)**, 284-304. (doi:10.2991/jnmp.2005.12.2.9)
- [15] Nucci MC, Leach PGL. 2005 Jacobi's last multiplier and the complete symmetry group of the Ermakov-Pinney equation. *J. Nonlinear Math. Phys.* **12(2)**, 305-320. (doi:10.2991/jnmp.2005.12.2.10)

- [16] Nucci MC, Leach PGL. 2007 Lagrangians galore. *J. Math. Phys.* **48**(12), 1–16, 123510. (doi: 10.1063/1.2821612)
- [17] Nucci MC, Leach PGL. 2008 Jacobi’s last multiplier and Lagrangians for multidimensional systems. *J. Math. Phys.* **49**(7), 1–8, 073517. (doi: 10.1063/1.2956486)
- [18] Nucci MC, Tamizhmani KM. 2010 Lagrangians for dissipative nonlinear oscillators: the method of Jacobi last multiplier. *J. Nonlinear Math. Phys.* **17**(02), 167–178. (doi:10.1142/S1402925110000696)
- [19] Gravel S, Winternitz P. 2002 Superintegrability with third-order integrals in quantum and classical mechanics. *J. Math. Phys.* **43**(12) 5902–5912. (doi:10.1063/1.1514385)
- [20] Rodriguez MA, Tempesta P, Winternitz P. 2009 Symmetry reduction and superintegrable Hamiltonian systems. *Journal of Physics: Conference Series*. **175**(1).
- [21] Winternitz P, Yurduen I. 2009 Integrable and superintegrable systems with spin in three-dimensional Euclidean space. *J. Phys. A: Math. Theor.* **42**(38) 385203. (doi:10.1088/1751-8113/42/38/385203)
- [22] Duarte LGS, Duarte SES, da Mota LACP, Skea JEF. 2001 Solving second-order ordinary differential equations by extending the Prelle-Singer method. *J. Phys. A: Math. Gen.* **34**(14), 3015. (doi:10.1088/0305-4470/34/14/308)
- [23] Chandrasekar VK, Senthilvelan M, Lakshmanan M. 2005 On the complete integrability and linearization of certain second-order nonlinear ordinary differential equations. *Proc. R. Soc. A*, **461**(2060), 2451–2477.
- [24] Chandrasekar VK, Pandey SN, Senthilvelan M, Lakshmanan M. 2006 A simple and unified approach to identify integrable nonlinear oscillators and systems. *J. Math. Phys.* **47**(2), 023508. (doi:10.1063/1.2171520)
- [25] Chandrasekar VK, Senthilvelan M, Lakshmanan M. 2006 A unification in the theory of linearization of second-order nonlinear ordinary differential equations. *J. Phys. A: Math. Gen.* **39**(3), L69. (doi:10.1088/0305-4470/39/3/L01)
- [26] Pradeep RG, Chandrasekar VK, Senthilvelan M, Lakshmanan M. 2010 On certain new integrable second order nonlinear differential equations and their connection with two dimensional Lotka–Volterra system. *J. Math. Phys.* **51**(3), 033519. (doi:10.1063/1.3327838)

- [27] Mohanasubha R, Chandrasekar VK, Senthilvelan M, Lakshmanan M. 2014 Interplay of symmetries, null forms, Darboux polynomials, integrating factors and Jacobi multipliers in integrable second-order differential equations. *Proc. R. Soc. A.* **470(2163)**, p. 20130656. (doi:10.1098/rspa.2013.0656)
- [28] Mohanasubha R, Chandrasekar VK, Senthilvelan M, Lakshmanan M. 2015 Interconnections between various analytic approaches applicable to third-order nonlinear differential equations. arXiv preprint arXiv:1502.03910.
- [29] Dimas S, Tsoubelis D. 2005 SYM: A new symmetry-finding package for Mathematica *Group Analysis of Differential Equations* Ibragimov NH, Sophocleous C & Damianou PA edd (University of Cyprus, Nicosia) 64-70. See also [http : // www.math.upatras.gr/ spawn](http://www.math.upatras.gr/spawn).
- [30] Dimas S, Tsoubelis D. 2006 A new Mathematica-based program for solving overdetermined systems of PDEs *8th International Mathematica Symposium* (Avignon, France).
- [31] Dimas S. 2008 *Partial Differential Equations, Algebraic Computing and Nonlinear Systems* (Thesis: University of Patras, Patras, Greece).
- [32] Andriopoulos K, Dimas S, Leach PGL, Tsoubelis D. 2009 On the systematic approach to the classification of differential equations by group theoretical methods *J. Comp. Appl. Math.* **230(1)**, 224-232. (doi:10.1016/j.cam.2008.11.002)
- [33] Polyanin AD, Zaitsev VF. 2003 *Handbook of Exact Solutions for Differential Equations*. Chapman & Hall/CRC, USA.
- [34] Tsuda T, Okamoto K, Sakai H. 2005 Folding transformations of the Painlevé equations. *Math. Ann.* **331(4)**, 713–738. (doi:10.1007/s00208-004-0600-8)
- [35] Grammaticos B, Ramani A, Willox R. 2011 Folding transformations and HKY mappings. *J. Nonlinear Math. Phys.* **18(01)**, 75–85. (doi: 10.1142/S1402925111001179)
- [36] Partha G, Choudhury AG. 2015 Folding transformations of equations from the Gambier family. *Commun. Nonlinear Sci. Numer. Simul.* **22(1)**, 1028–1035. (doi:10.1016/j.cnsns.2014.09.021)